

Evaluation of Two Nonradiolabeled Murine Local Lymph Node Assays (LLNA) for Potency Categorization of Substances Causing Allergic Contact Dermatitis in Humans

J Strickland¹, F Stack¹, T Burns¹, D Allen¹, W Casey², W Stokes²
¹ILS, Inc., RTP, NC, USA; ²NICEATM/NTP/HHS, RTP, NC, USA

Abstract

The correct classification of strong skin sensitizers is critical because such substances are considered to have a significant potential for causing allergic contact dermatitis (ACD) in humans. Because the prognosis for ACD is poor, sensitizing substances must be labeled with a description of the potential hazard and the precautions necessary for workers and consumers to avoid development of ACD. A recent ICCVAM evaluation found that the LLNA correctly classified 52% (14/27) of the strong human sensitizers when an effective threshold concentration (EC) $\leq 2\%$ was used as the criterion. Thus, ICCVAM recommends that the LLNA may be used as a screening test to classify substances as strong sensitizers, but that the classification of substances as other than strong sensitizers requires additional information. The OECD recently adopted two test guidelines for nonradiolabeled versions of the LLNA that could be used to classify substances as sensitizers: the LLNA: BrdU-ELISA and the LLNA: DA. Although these LLNA methods use different decision criteria to classify substances for ACD hazard, their accuracy is comparable to that of the LLNA. Of the 136 substances used in the ICCVAM evaluation of the usefulness of the LLNA for potency categorization, LLNA: BrdU-ELISA data were available for 31 substances, and LLNA: DA data were available for 30 substances. An EC $\leq 9\%$ for the LLNA: BrdU-ELISA and an EC $\leq 0.8\%$ for the LLNA: DA classified strong human sensitizers at rates comparable to that of the LLNA. These results suggest that the LLNA: BrdU-ELISA and the LLNA: DA may also be useful for classifying substances as strong human sensitizers.

Introduction

- Allergic contact dermatitis (ACD) is a skin reaction characterized by redness, swelling, and itching that can result from repeated contact with a sensitizing substance.
- Because the prognosis is poor (Hogan et al. 1990), preventing or limiting exposures to ACD hazards is important.



- The U.S. Consumer Product Safety Commission (CPSC), under the Federal Hazardous Substances Act (15 U.S.C. 1261-1278), requires hazard labeling of strong skin sensitizers (16 CFR 1500.3(b)(9)).
- Although the CPSC uses both human and animal data to determine that a substance is a strong sensitizer, no quantitative criteria are currently applied (Table 1).
- In 2007 the CPSC requested that ICCVAM evaluate the usefulness and limitations of the LLNA as a stand-alone test method for potency determination.
- Following the CPSC request, the Globally Harmonized System for Classification and Labeling of Chemicals (GHS) was updated to include potency criteria for the LLNA (UN 2009, 2011) (Table 1).
- Based on an evaluation of 136 substances with LLNA and human data, ICCVAM recommended that LLNA with an EC $\leq 2\%$ could be used to classify substances as strong sensitizers (GHS 1A) but that LLNA with EC $> 2\%$ could not be used alone to classify substances as other (than strong) sensitizers (GHS 1B) (ICCVAM 2011).
- 48% (13/27) of strong human sensitizers were classified as other sensitizers (EC $> 2\%$) or as nonsensitizers by the LLNA.
- This poster examines the accuracy of two nonradiolabeled versions of the LLNA, the LLNA: BrdU-ELISA and the LLNA: DA, for classifying substances as strong human sensitizers.

Table 1. Classification Systems for Skin Sensitizers


Classification System	Criteria for "Strong" Skin Sensitizer	Criteria for "Other" (Than Strong) Skin Sensitizer
CPSC	Weight-of-evidence approach that considers quantitative or qualitative risk assessment, frequency of occurrence and range of severity of reactions in healthy or susceptible populations, and the results of experimental assays in animals or humans. Human data take precedence over animal data, other data on potency or bioavailability of sensitizers, data on reactions to a cross-reacting substance, or to a chemical that metabolizes or degrades to form the same or a cross-reactive substance, the threshold of human sensitivity, epidemiological studies, case histories, occupational studies, and other appropriate <i>in vivo</i> and <i>in vitro</i> test studies.	Not applicable
GHS	Subcategory 1A High frequency of occurrence in humans, and/or high potency in animals. May consider severity. LLNA EC3 $\leq 2\%$ Positive ¹ response in humans at $\leq 500 \mu\text{g}/\text{cm}^2$	Subcategory 1B Low to moderate frequency of occurrence in humans and/or low to moderate potency in animals. LLNA EC3 $> 2\%$ Positive ² response in humans at $> 500 \mu\text{g}/\text{cm}^2$

Abbreviations: CPSC = U.S. Consumer Product Safety Commission; EC3 = estimated concentration of a substance needed to produce a stimulation index of 3 in the LLNA; GHS = Globally Harmonized System for Classification and Labeling of Chemicals; LLNA = murine local lymph node assay.
¹ Human evidence can also include (1) diagnostic patch test data with a relatively high incidence of reactions in a defined population in relation to relatively low exposure or (2) other epidemiologic evidence with a relatively high incidence of allergic contact dermatitis in relation to relatively low exposure.
² Human evidence can also include (1) diagnostic patch test data with a relatively low but substantial incidence of reactions in a defined population in relation to relatively high exposure or (2) other epidemiologic evidence with a relatively low but substantial incidence of allergic contact dermatitis in relation to relatively high exposure.

Methods


- LLNA: BrdU-ELISA Test Method**
 - The assay measures the nucleotide analog bromodeoxyuridine (BrdU) using an enzyme-linked immunosorbent assay (ELISA) to assess lymph node cell proliferation (Takeyoshi et al. 2001) (Figure 1).
 - ICCVAM recommended use of the LLNA: BrdU-ELISA to classify substances as potential sensitizers (stimulation index [SI] ≥ 1.6) or nonsensitizers (ICCVAM 2010a).
 - OECD Test Guideline 442B Skin Sensitization: Local Lymph Node Assay: BrdU-ELISA, which includes the SI ≥ 1.6 criterion to classify substances as skin sensitizers, was adopted on July 22, 2010 (OECD 2010a).
- LLNA: DA Test Method (developed by Daiichi Chemical Industries, Ltd.)**
 - The assay measures adenosine triphosphate content in draining auricular lymph nodes as an estimate of cell number to assess lymph node cell proliferation (Yamashita et al. 2005; Idehara et al. 2008) (Figure 2).
 - ICCVAM recommended use of the LLNA: DA to classify substances as potential sensitizers (SI ≥ 1.8) or nonsensitizers (ICCVAM 2010b).
 - OECD Test Guideline 442A Skin Sensitization: Local Lymph Node Assay: DA, which includes the SI ≥ 1.8 criterion to classify substances as skin sensitizers, was adopted on July 22, 2010 (OECD 2010b).

Figure 1. LLNA: BrdU-ELISA Test Method Protocol



Abbreviations: SI = stimulation index.

Figure 2. LLNA: DA Test Method Protocol



Abbreviations: RLU = relative luminescence unit; SI = stimulation index; SLS = sodium lauryl sulfate.

Human Data

- These data were obtained from the literature as reported by ICCVAM (2011).
- The human repeat insult patch test (HR IPT) and the human maximization test (HMT) involve the administration of occluded patches loaded with test substance to the skin for 0 (HR IPT) or 5 (HMT) on-and-off periods of 24 or 48 hours, respectively, in order to attempt to induce an allergic reaction (Kligman and Epstein 1975; Politano and Api 2008).
- Following a rest period of 10 to 14 days, volunteers are again exposed to the test substance in an occluded patch on naive skin for 24 (HR IPT) or 48 (HMT) hours.
- Erythema and edema (including papules, vesicles, or bullae) observed after patch removal indicate ACD and are recorded as positive reactions.
- For substances that produce no skin irritation, the HMT includes a patch pretreatment of the skin with 5% sodium lauryl sulfate for the 24-hour period before the induction patch treatments in order to break the stratum corneum barrier, producing a "brisk dermatitis" (Kligman and Epstein 1975).
- Induction thresholds (the minimum concentrations that produce positive reactions) are reported as microgram weight of applied substance per square centimeter of skin ($\mu\text{g}/\text{cm}^2$).

Chemical Database for Analysis

- Data for analysis were identified from the corresponding ICCVAM evaluations of the LLNA: BrdU-ELISA (ICCVAM 2010a), the LLNA: DA (ICCVAM 2010b), and the traditional LLNA for potency categorization (ICCVAM 2011) (see Table 2) and included:
 - 30 substances with traditional LLNA, human, and LLNA: BrdU-ELISA data
 - 31 substances with traditional LLNA, human, and LLNA: DA data
- Human studies were used as the reference data.
 - Induction threshold concentrations from the HR IPT and HMT were expressed as the dose per unit area that produced a 5% response in the population tested (DSA_{50}).
 - "Strong" sensitizers (GHS 1A) are defined as having $\text{DSA}_{50} \leq 500 \mu\text{g}/\text{cm}^2$. "Other" sensitizers (GHS 1B) are defined as having $\text{DSA}_{50} > 500 \mu\text{g}/\text{cm}^2$.

Analyses

- The correct, underprediction, and overprediction rates for strong (GHS 1A) human sensitizers, other (GHS 1B) human sensitizers, and nonsensitizers were calculated for multiple threshold induction concentrations for the LLNA: BrdU-ELISA EC1.6 and the LLNA: DA EC1.8.
- These results were compared to the prediction rates of the traditional LLNA EC3 for the same substances.

*EC1.6 is an estimated concentration of a substance needed to produce a stimulation index of 1.6 in LLNA: BrdU-ELISA.
 *EC1.8 is an estimated concentration of a substance needed to produce a stimulation index of 1.8 in LLNA: DA.

Acknowledgements

The Intramural Research Program of the National Institute of Environmental Health Sciences (NIEHS) supported this poster. Technical support was provided by ILS, Inc., under NIEHS contract N01-ES-35504. This poster reflects the views of the authors. The views expressed above have not been reviewed or approved by any U.S. Federal agency and do not necessarily represent the official positions of any U.S. Federal agency.

Because the poster was written as part of the official duties of the authors, it can be freely copied.

NICEATM gratefully acknowledges the following individuals and institutions for submitting data to NICEATM for the LLNA: BrdU-ELISA and LLNA: DA evaluations.

- | | |
|--|---|
| Kenji Idehara, PhD
Daiichi Chemical Industries, Ltd.
Osaka, Japan | Takashi Omori, PhD
Kyoto University School of Public Health
Kyoto, Japan |
| Masahiro Takeyoshi, PhD
Japanese Center and Research Institute
Saitama, Japan | Hajime Kojima, PhD
Japanese Center for the Validation of Alternative Methods
National Institute of Health Sciences
Ministry of Health, Labour and Welfare
Tokyo, Japan |


Posters and abstracts presented by NICEATM-ICCVAM at SOT 2012 are available on the NICEATM-ICCVAM website at <http://iccvam.niehs.nih.gov/>.

Table 2. Database for Potency Analysis of Nonradiolabeled LLNA Methods

Substance ^a	Human DSA_{50} ($\mu\text{g}/\text{cm}^2$) ^b	LLNA EC3 ^c	LLNA: BrdU-ELISA EC1.6 ^d	LLNA: DA EC1.8 ^e
2,4-Dinitrochlorobenzene	3.4	0.04	0.02	0.02
5-Chloro-2-methyl-4-isothiazolin-3-one	5.0	0.01	0.07	0.01
p-Phenylenediamine	30	0.12	NC	0.04
Potassium dichromate	106	0.12	NT	0.06
Formaldehyde	191	1.40	0.08	0.47
Cobalt chloride	279	0.57	0.32	0.38
Cinnamic aldehyde	382	1.00	4.81	0.63
Diethyl malate	400	3.27	8.05	0.89
Butyl glycidyl ether	437	30.9	NT	17.5
Ethyl acrylate	818	32.0	33.3	6.9
Citral	915	5.00	NC	4.11
Isounginol	1016	1.40	4.93	0.62
Quararibiphenyl	1073	0.16	0.08	0.1
2-Mercaptobenzothiazole	1930	2.40	12.1	1.99
Aniline	2463	33.0	13.6	NT
Cinnamic alcohol	3002	20.0	74	5.23
Imidazolidinyl urea	3846	24.0	49.6	6.28
Sulfanilamide	4310	NC	NC	NC
Hydroxytolonil	5327	23.0	17.1	8.67
Eugenol	5926	11.0	6.8	2.63
Benzocaine	10140	78.0	NT	3.11
Phenyl benzoate	52500	9.50	16.9	0.65
Benzalkonium chloride	Negative	0.07	NT	0.4
Cyclamen aldehyde	Negative	22.0	28.9	NT
Diethyl phthalate	Negative	NC	NC	NC
Cyclozol	Negative	NC	NC	NT
Hioxane	Negative	NC	NC	82.2
Hexyl chromic aldehyde	Negative	8.90	10.2	6.16
Isopropenol	Negative	NC	5.33	NC
Isopropyl myristate	Negative	44.0	NC	NT
Linalool	Negative	55.0	27.6	NT
Methyl salicylate	Negative	17.0	NC	NC
Propylene glycol	Negative	NC	NC	NT
Resorcinol	Negative	5.9	NT	3.9
Salicylic acid	Negative	12.0	NC	17.7
Sodium lauryl sulfate	Negative	4.0	13.3	1.64

Abbreviations: DSA_{50} = induction dose per skin area, in $\mu\text{g}/\text{cm}^2$, in a human repeat insult patch test or human maximization test that produces a positive response in 5% of the tested population; EC1.6 = estimated concentration of a substance needed to produce a stimulation index of 1.6 in the LLNA: BrdU-ELISA; EC1.8 = estimated concentration of a substance needed to produce a stimulation index of 1.8 in the LLNA: DA; EC3 = estimated concentration of a substance needed to produce a stimulation index of 3 in the traditional LLNA; NC = not calculated because the substance was negative; NT = not tested.
^aListed in order of decreasing human potency.
^bMore than one value per substance was identified; values represent a geometric mean.

Figure 3. LLNA: BrdU-ELISA and Human Potency



Abbreviations: DSA_{50} = induction dose per skin area, in $\mu\text{g}/\text{cm}^2$, in a human repeat insult patch test or human maximization test that produces a positive response in 5% of the tested population; EC1.6 = estimated concentration needed to produce a stimulation index of 1.6 in the LLNA: BrdU-ELISA.
 Notes: A red vertical line indicates a classification cutoff of 9% for the LLNA: BrdU-ELISA EC1.6. A red horizontal line shows a cutoff of 500 $\mu\text{g}/\text{cm}^2$ for the human DSA_{50} — a cutoff applied by GHS to differentiate "strong" sensitizers from "other" sensitizers. Negative results were assigned a single value $> 100,000 \mu\text{g}/\text{cm}^2$ for the human DSA_{50} . Six true negatives are overlaid in the unshaded area in the upper right corner of the graph, and the number of false positives and false negatives (relative to human data) are indicated on the graph.

Results

Figures 3 and 4 show scatter plots for the human and LLNA: BrdU-ELISA or LLNA: DA data in Table 2. Table 3 shows classification rates for the nonradiolabeled LLNA assays to predict human potency based on the GHS cutoff of 500 $\mu\text{g}/\text{cm}^2$ for the human DSA_{50} . The two figures and table are color coded to indicate substances that are correctly classified (unshaded), overclassified (yellow), or underclassified (pink) using selected cutoffs for the two LLNA tests.

Table 3. Classification Rates for Prediction of Human Potency

Classification Cutoff for "Strong" versus "Other" Sensitizers	"Strong" Human Sensitizers ($\text{DSA}_{50} \leq 500 \mu\text{g}/\text{cm}^2$)		"Other" Human Sensitizers ($\text{DSA}_{50} > 500 \mu\text{g}/\text{cm}^2$)		Human Nonsensitizers		Overall Correct Potency Classification ^a	
	Correct	Under	Under	Correct	Over	Over		
LLNA EC3 $\leq 2\%$	86%	14%	8%	75%	17%	58%	42%	65%
LLNA: BrdU-ELISA EC1.6 $\leq 9\%$	86%	14%	17%	54%	25%	50%	50%	61%
LLNA: BrdU-ELISA EC1.6 $\leq 6\%$	71%	29%	17%	67%	17%	50%	50%	61%
LLNA: BrdU-ELISA EC1.6 $\leq 2\%$	57%	43%	17%	75%	28%	50%	50%	61%


Abbreviations: DSA_{50} = induction dose per skin area, in $\mu\text{g}/\text{cm}^2$, in a human repeat insult patch test or human maximization test that produces a positive response in 5% of the tested population; EC1.6 = estimated concentration of a substance needed to produce a stimulation index of 1.6 in the LLNA: BrdU-ELISA; EC3 = estimated concentration of a substance needed to produce a stimulation index of 3 in the traditional LLNA; EC3 = estimated concentration of a substance needed to produce a stimulation index of 3 in the LLNA: DA; EC3 = estimated concentration of a substance needed to produce a stimulation index of 3 in the LLNA: DA.
 Note: Column headings are defined as follows: correct = correct classification; under = underclassification; over = overclassification.
^aPotency classification used cutoff values of $\leq 2\%$, 6%, or 9% for strong sensitizers and $> 2\%$, 6%, or 9% for other sensitizers. "Strong" or "other" sensitizers were identified using the respective EC values listed in the table for the traditional LLNA and nonradiolabeled LLNA assays. The cutoff for the LLNA: BrdU-ELISA is indicated in the table (see row with the bold header for EC1.6 $\leq 9\%$).
 Note: The overall correct classification rate is based on human data classifications for "strong" sensitizers, "other" sensitizers, and nonsensitizers (UN 2011).

Table 3b. LLNA EC3 and LLNA: DA EC1.8 at Various Cutoffs^a for 30 Substances

Classification Cutoff for "Strong" versus "Other" Sensitizers	"Strong" Human Sensitizers ($\text{DSA}_{50} \leq 500 \mu\text{g}/\text{cm}^2$)		"Other" Human Sensitizers ($\text{DSA}_{50} > 500 \mu\text{g}/\text{cm}^2$)		Human Nonsensitizers		Overall Correct Potency Classification ^a	
	Correct	Under	Under	Correct	Over	Over		
LLNA EC3 $\leq 2\%$	86%	14%	8%	75%	17%	58%	42%	65%
LLNA: DA EC1.8 $\leq 2\%$	86%	14%	17%	54%	25%	50%	50%	61%
LLNA: DA EC1.8 $\leq 0.8\%$	76%	24%	17%	54%	17%	47%	33%	43%
LLNA: DA EC1.8 $\leq 0.5\%$	67%	33%	17%	47%	17%	33%	23%	43%

Abbreviations: See Table 3a.
 Note: Column headings are defined as follows: correct = correct classification; under = underclassification; over = overclassification.
^aPotency classification used cutoff values of $\leq 0.5\%$, 0.8%, or 2% for "strong" sensitizers and $> 0.5\%$, 0.8%, or 2% for "other" sensitizers. Sensitizers were identified using the respective EC values listed in the table for the traditional LLNA and the two nonradiolabeled LLNA assays. The cutoff for the LLNA: DA is indicated in the table (see row with bold header for EC1.8 $\leq 0.8\%$).
 Note: The overall correct classification rate is based on human data classifications for "strong" sensitizers, "other" sensitizers, and nonsensitizers (UN 2011).


Figure 4. LLNA: DA and Human Potency



Abbreviations: DSA_{50} = induction dose per skin area, in $\mu\text{g}/\text{cm}^2$, in a human repeat insult patch test or human maximization test that produces a positive response in 5% of the tested population; EC1.8 = estimated concentration needed to produce a stimulation index of 1.8 in the LLNA: DA.
 Notes: A red vertical line indicates a classification cutoff of 0.8% for the LLNA: DA EC1.8. A red horizontal line shows a cutoff of 500 $\mu\text{g}/\text{cm}^2$ for the human DSA_{50} — a cutoff applied by GHS to differentiate "strong" sensitizers from "other" sensitizers. Three true negatives are overlaid in the unshaded area in the upper right corner of the graph, and the number of false positives and false negatives (relative to human data) are indicated on the graph.


Figure 5. Classification of "Strong" Sensitizers

Figure 5a. Cutoffs for the LLNA: BrdU-ELISA EC1.6



Abbreviations: EC1.6 = estimated concentration of a substance needed to produce a stimulation index of 1.6 in the LLNA: BrdU-ELISA; EC3 = estimated concentration of a substance needed to produce a stimulation index of 3 in the LLNA.
 Note: Cutoffs are compared using values from Table 3a and Figure 3.

Figure 5b. Cutoffs for the LLNA: DA EC1.8



Abbreviations: EC1.8 = estimated concentration of a substance needed to produce a stimulation index of 1.8 in the LLNA: DA; EC3 = estimated concentration of a substance needed to produce a stimulation index of 3 in the LLNA.
 Note: Cutoffs are compared using values from Table 3b and Figure 4.

Conclusions

- EC1.6 $\leq 9\%$ for the LLNA: BrdU-ELISA and EC1.8 $\leq 0.8\%$ for the LLNA: DA classified strong sensitizers at rates comparable to that of the LLNA EC3 $\leq 2\%$.
- These results suggest that the LLNA: BrdU-ELISA and LLNA: DA may also be useful for classifying substances as strong sensitizers.
- When relative potency information is required, these nonradiolabeled tests should
 - Further reduce and refine animal use for ACD hazard assessments in comparison to guinea pig test methods, while ensuring human safety
 - Reduce costs and environmental hazards associated with the use of radioactive substances

References

Hogan DJ, Dannaker CJ, Malbach H. 1990. The prognosis of contact dermatitis. *J Am Acad Dermatol* 23:300-30.

ICCVAM. 2010a. ICCVAM Test Method Evaluation Report on the Murine Local Lymph Node Assay: BrdU-ELISA. A Nonradioactive Alternative Test Method to Assess the Allergic Contact Dermatitis Potential of Chemicals and Products. NIH Publication No. 10-7552. Research Triangle Park, NC: National Institute of Environmental Health Sciences. Available: <http://iccvam.niehs.nih.gov/methods/immunotox/lna-ELISATMER.htm>

ICCVAM. 2010b. ICCVAM Test Method Evaluation Report on the Murine Local Lymph Node Assay: DA. A Nonradioactive Alternative Test Method to Assess the Allergic Contact Dermatitis Potential of Chemicals and Products. NIH Publication No. 10-7551. Research Triangle Park, NC: National Institute of Environmental Health Sciences. Available: <http://iccvam.niehs.nih.gov/methods/immunotox/lna-DATMER.htm>

ICCVAM. 2011. ICCVAM Test Method Evaluation Report: Usefulness and Limitations of the Murine Local Lymph Node Assay for Potency Categorization of Chemicals Causing Allergic Contact Dermatitis in Humans. NIH Publication No. 11-7709. Research Triangle Park, NC: National Institute of Environmental Health Sciences. Available: <http://iccvam.niehs.nih.gov/methods/immunotox/lna-potTMR.htm>

Idehara K, Yamagishi G, Yamashita K, Ito M. 2008. Characterization and evaluation of a modified local lymph node assay using ATP content as a non-radio isotopic endpoint. *J Pharmacol Toxicol Meth* 58: 1-10.

Kligman AM, Epstein WI. 1975. Updating the maximization test for identifying contact allergens. *Contact Dermatitis* 1(4): 231-238.

OECD. 2010a. Test No. 442B. Skin Sensitization: Local Lymph Node Assay: BrdU-ELISA [adopted 22 July 2010]. In: OECD Guidelines for the Testing of Chemicals, Section 4: Health Effects. Paris: OECD Publishing. Available: http://www.oecd-ilibrary.org/environnement/test-no-442b-skin-sensitization_3788264909972-en

OECD. 2010b. Test No. 442A. Skin Sensitization: Local Lymph Node Assay: DA [adopted 22 July 2010]. In: OECD Guidelines for the Testing of Chemicals, Section 4: Health Effects. Paris: OECD Publishing. Available: